

## **Participant Initial Questionnaire**

Study title: Reflex testing for MAFLD in patients with type 2 diabetes

Chief Investigator: Professor Christopher Byrne Participant Identification Number: \_\_\_\_\_\_\_

Date: Researcher:							
Date of Birth (day/month/year)							
Sex							
Ethnicity							
Current prescription medications							
AUDIT-C <sup>1</sup> Bush K, Kivlahan DR, et al (1998). The AUDIT alcohol consumption questions (AUDIT-C): An effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Arch Intern Med. 158:1789-95)							
How often do you have a drink containing alcohol?	Never	Monthly or less	2-4 tin month		2-3 times a week	4 or more times a week	
How many standard drinks containing alcohol do you have on a typical day?	1 or 2	3 to 4	5 to 6		7 to 9	10 or more	
How often do you have six or more drinks on one occasion?	Daily or almost daily	Weekly	Month	nly	Less than monthly	Never	
AUDIT-C score							
Venesection performed?	Yes / No						
Location and number of attempts							
Blood samples obtained?	Yes / No						
Refer patient elsewhere for venesection	Yes / No If yes, remember to provide patient with labels for test tubes						
Liver Assessment							
Has the patient fasted? Y/N	Date:			Probe	size: XL / M		



VCTE reading (kPa):	IQR/MED:	CAP (Db/m²):			
Patient advised of VCTE assessment? Y/N					
Additional information					